

Quantum.Silver vir Pluimvee – Antimikrobies

Reg Nr: Act 36/1947 V 35062

'n Nuwe generasie produk, wat Quantum Dot-tegnologie gebruik om 'n natuurlike silwerproduk met mikro-dodende effekte te vervaardig. Quantum.Silver® help met die behandeling van infeksies: Virusse, Bakterieël, Swam- en Koksidiale infeksies

Verpakking:

Oplossing: 1, 5 en 25 Liter

Voormengsel: 5kg emmer

Dosering: Oplossing: 1 ml/kg liggaamsmassa daaglik vir 5-14 dae
of meng 25 ml/1 L drinkwater

Voormengsel: QS Premix teen 200g/ton tot 400g/ton vir 1-2 weke.

Roetinebehandelings:

1. Broilers – Dag 1-3 in drinkwater (1% QS in drinkwater)
2. Broilers – Dag 21-24 in drinkwater (1% QS in drinkwater)
3. Broilers – 3 dae voor slagting in drinkwater
4. Broilers – as deel van 'n skoffelprogram saam met ander anti-koksidiale middels
5. Lèhen: Behandel volwasse hen vir 14 dae na aankoms op die plaas (400g/ton premix of in water)
6. Lèhen : Behandel volwasse hen elke 2 of 3 maande vir 14 dae in die voer met 400g/ton premix

Onttrekkingsperiode:

As gevolg van die baie nano-grootte van die silwerdeeltjies, word dit vinnig uitgeskei en het dus geen vleis- of eieronttrekkingsperiode nie.


Inentings: Skakel Quantum.Silver-behandeling uit vir 1 dag voor inenting en tot 2 dae na lewende virusinentings.

Antibiotika:

Quantum.Silver kan saam met antibiotika of enige ander medikasie gebruik word.

Nuwe effekte: Daar is geen bekende nuwe effekte van Quantum.Silver nie.

Vir meer inligting, kontak: Dr. Thomas Volker 084 992 2194 of thomas@eaglec.co.za



Quantum.Silver for Poultry – Antimicrobial

Reg Nr: Act 36/1947 V 35062

A new generation product, using the Quantum Dot technology to produce a natural silver product with micro-cidal effects, which aids in the treatment of infections:

1. Viral, Bacterial, Fungal and Coccidial

Packaging:

Solution: 1, 5 and 25 Liter

Premix: 5kg bucket

Dosage: Solution: 1 ml/kg body mass daily for 5- 14 days

Mix 25 ml/1 L of drinking water

Premix: QS Premix at 200g/ton to **400g/ton** for 1- 2 weeks.

Recommended routine treatments:

1. Broiler chicks – Day 1- 3 in drinking water (1% QS in drinking water)
2. Broiler chicks – Day 21 -24 in drinking water(1% QS in Drinking water)
3. Broiler chicks – 3 days prior to slaughter in drinking water
4. Broiler chicks - as part of a shuttle program with other anti-coccidials
5. Layers: treat hens for 7 days after arrival on farm (200g/ton premix or in water)
6. Layers: treat hens every 2 or 3 months for 14 days in feed with 400g/ton Premix

Withdrawal period:

Due to the very nano- size of the silver particles, it is excreted quickly and therefore has no meat or egg withdrawal period.

Vaccinations: Withdraw Quantum.Silver treatment for 1 day prior to vaccination till 2 days post live viral vaccinations.

Antibiotics:

Quantum.Silver may be used in conjunction with antibiotics or any other medication.

Side effects: There are no known side effects for Quantum.Silver

For more info, contact: Dr Thomas Volker 084 992 2194 or thomas@eaglec.co.za